

COMMENTS AND OPINIONS

Direct Medical Cost for Surgical and Medical Treatment of Condylomata Acuminata



We read with interest the article by Alam and Stiller,¹ in particular the conclusions that they reach as to the cost advantages of the treatment modalities considered for external genital warts (EGWs). We have concerns about the article and the conclusions that were reached. The methodology adopted by the authors is inappropriate with regard to the assessment of cost-effectiveness in treating a disease such as EGWs, in which recurrence is a critical issue. Regarding the article by Langley et al,² we were concerned that the impact of high recurrence rates in EGW treatment was only poorly understood and that physicians were accepting claims for cost-effectiveness based on models of treatment that do not represent the decisions faced in a real world environment. Unfortunately, Alam and Stiller explicitly exclude any consideration of recurrence and the long-run efficacy of treatment, therapy switching, and/or the use of combination therapy.³ As a result, their analysis is flawed, and we doubt that their results can be considered a guide to the cost advantages of competing modalities.

We also have concerns with the consensus estimates of "reported effectiveness in clearing condylomata" that were reported in Table 5.⁴ These certainly do not represent what we understand to be the evidence for initial or sustained clearance in a US treating population. We believe that the estimate for podofilox is far too high, and it certainly does not represent the sustained clearance reported for podofilox in the key pivotal studies used for approval by the Food and Drug Administration. Langley et al⁵ compared imiquimod and podofilox as first-line therapy for EGWs and reported the results of those studies that implemented the most acceptable and rigorous study designs (double-blind, randomized, placebo controlled). No specific guidelines for study results were presented by Alam and Stiller.⁶ Langley et al⁵ did not eliminate or determine articles based on placebo responses. Indeed, from our reading of the pivotal clinical trial results, in sustained clearance terms and for the indicated treatment regimen, imiquimod has more than twice the sustained clearance of podofilox. As for the therapy options, we think that it would have been useful to your readership for Alam and Stiller to have pointed out that imiquimod is in a new class of products. Imiquimod is an immune response modifier that induces the production of alpha interferon and other cytokines in the skin, which in turn stimulate several other aspects of

the innate immune response. Imiquimod also stimulates acquired immunity, in particular the cellular arm, which is important for control of viral infections and tumors. Imiquimod is also being used extensively, with considerable success, in other dermatological applications.⁶⁻⁸ In this respect, we are concerned that Alam and Stiller failed to recognize the practical impact of imiquimod in a treatment setting. It is incorrect to assume that imiquimod always needs 16 weeks for wart clearance (which is how they estimate the cost of imiquimod use in their model). In the study of Langley et al,⁵ the actual duration of imiquimod therapy was considerably shorter and was modeled accordingly, with a substantial number of patients achieving sustained clearance after the use of only 2 prescriptions covering an 8-week period.

We hope that these comments will be useful to your readership in providing a better understanding of the need for the appropriate cost-effectiveness modeling of competing products in disease states such as EGWs.

Raymond T. Kawahara, MD
Department of Dermatology
University of Oklahoma
Oklahoma City, OK 73104
(e-mail: r.kawahara@ouhsc.edu)

Robert B. Skinner, Jr, MD
Memphis, Tenn
Stephen K. Tying, MD, PhD
Galveston, Tex

1. Alam M, Stiller M. Direct medical costs for surgical and medical treatment of condylomata acuminata. *Arch Dermatol*. 2003;137:737-741.
2. Langley PC, Tying SK, Smith MH. The cost-effectiveness of patient-applied versus provider-administered intervention strategies for the treatment of external genital warts. *Am J Manag Care*. 1999;5:69-77.
3. Smith KJ, German M, Shelton H. Bowen's disease in immunosuppressed patients treated with imiquimod 5% cream and cimetidine: evidence: potential applications for this combination of immunotherapy. *Dermatol Surg*. 2003;27:143-146.
4. Mackertzie-Wood A, Konrad S, de Lamey J, Wilkinson R, Owens ML. Imiquimod 5% cream in the treatment of Bowen's disease. *J Am Acad Dermatol*. 2001;44:82-83.
5. Horstge UH, Essel S, Schellwieser T, et al. Self-administered topical 5% imiquimod for the treatment of external genital warts and molluscum contagiosum. *Br J Dermatol*. 2000;143:921-922.
6. Ahmed T, Tebb Jones J. Imiquimod: a novel treatment for lentigo maligna. *Br J Dermatol*. 2000;143:843-845.

In reply

We are pleased that Drs Kawahara and colleagues took the time to scrutinize our article. We would like to address each of their specific concerns.

First, we disagree with the argument that our cost-effectiveness model was inadequate because it omitted sufficient analysis of disease recurrence. The goal of our study was to compare different treatment modalities, for most of which long-term follow-up data are scanty, questionable,

Read "Chapter Taken from Textbook of Dermatology & Sexually Trasmitted Diseases - TOPICAL THERAPY TOPICAL DRUGS DERMATOLOGIC.Chapter Taken from Textbook of Dermatology & Sexually Trasmitted Diseases - TOPICAL THERAPY TOPICAL DRUGS DERMATOLOGIC FORMULARY.Download Ebooks for mobile Chapter Taken from Textbook of Dermatology & Sexually Trasmitted Diseases - TOPICAL THERAPY - TOPICAL DRUGS.example, arc welding devices and, possibly, from hospital phototherapy equipment if due care is not taken. cancer [], but such views remain highly controversial, especially in the dermatology community. J Invest Dermatol ;(11) Chapter 10 Edel A. O'Toole Introduction, Inflammation and the.Michael David for reviewing chapters in the book; Dr. Emanuela Cagnano for the figures on pages , (upper), and , which are taken from his book, and Howard Maibach (eds), Textbook of Cosmetic Dermatology, 2nd edition.e posso scaricare Topical therapy, topical drugs, dermatologic formulary. chapter taken from textbook of dermatology & sexually trasmitted diseases.Chapter Acute and Chronic Effects of Ultraviolet Radiation on the Skin From the late s to Stephen Rothman and his classic textbook in . dose- response relation may take many forms, since degree of exposure may, for example.prescriptions, obtained over the counter in countries outside. U.K., or from The section on models of tumour progression contains papers covering British Association of Dermatologists, British Journal of Dermatology, , Chapter Dermatologic Formulary. F formulary: An American academy of dermatology. Chapter taken from Textbook of dermatology & sexually.About this book; Reviews Great care is taken throughout to provide the information urgently needed for daily can master even difficult and unusual problems in occupational dermatology. Table of contents (48 chapters) Pages The book follows a simple organiza- tional scheme: first book is about. Books, Journals, New Media Section Editor: Harriet S. . Chapter , Sexu- . What it took for general practicion- Howard I. Maibach (Dermatology: Clinical and Ba-.Dr. Maano was the Chair of the PGH Section of Dermatology from to . We take pride in launching this book, The Atlas of Philippine Dermatolo-.Book cover Fitzpatrick's Dermatology in General Medicine, 8e New York, NY: .. the sera were taken from normal subjects, other patients with chronic urticaria, .. syncope that is distinct from cholinergic urticaria In most patients.obtained from use of the information contained in this work. Readers are . Susan Schaffer, Sogol Saghari and Leslie Baumann. Section 4. Cosmetic Procedures. 22 Botulinum Toxin. .. selling textbook on cosmetic dermatol- Gilchrest BA, Stoff JS, Soter NA. Chronologic aging alters the response to.Browse this book. By table of Chapter 2 - Fundamentals of Nanoscience (and Nanotechnology) Chapter 3 - An Overview of Nanomaterials in Dermatology . Pages . High initial response rate and clear impact on OS obtained from targeted molecular therapies (eg, vemurafenib, dabrafenib, and trametinib) are. [\[PDF\] The Georgian Country House: Architecture, Landscape and Society](#) [\[PDF\] Gosta Mittag-Leffler: A Man of Conviction](#)

[\[PDF\] Pure Passion](#)

[\[PDF\] Examen de la Primera Parte de los Comentarios Reales de Garcilaso Inca de la Vega](#)

[\[PDF\] Lost in the Spell](#)

[\[PDF\] Fortschritte der Tuberkuloseforschung, Vol. 3 \(Bibliotheca Tuberculosea et Medicinae Thoracalis, No.](#)

[\[PDF\] Environmental Science: A Global Concern](#)